International
Scientific Conference

Algebraic and Geometric Methods of Analysis



26-30 may 2020
Odesa, Ukraine

List of topics

- Algebraic methods in geometry
- Differential geometry in the large
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Dynamical systems and their applications
- Geometric problems in mathematical analysis
- Geometric and topological methods in natural sciences

Organizers

- Ministry of Education and Science of Ukraine
- Odesa National Academy of Food Technologies
- Institute of Mathematics of the National Academy of Sciences of Ukraine
- Odessa I. I. Mechnikov National University
- Taras Shevchenko National University of Kyiv
- International Geometry Center
- Kyiv Mathematical Society

Program Committee

Chairman: Prishlyak A.
(*Kyiv, Ukraine*)
Balan V.
(*Bucharest, Romania*)
Banakh T.
(*Lviv, Ukraine*)
Bolotov D.
(*Kharkiv, Ukraine*)
Borysenko O.
(*Kharkiv, Ukraine*)
Cherevko Ye.
(*Odesa, Ukraine*)
Fedchenko Yu.
(*Odesa, Ukraine*)
Karlova O.
(*Chernivtsi, Ukraine*)
Kiosak V.
(*Odessa, Ukraine*)
Kirillov V.
(*Odesa, Ukraine*)
Konovenko N.
(*Odesa, Ukraine*)
Lyubashenko V.
(*Kyiv, Ukraine*)
Maksymenko S.
(*Kyiv, Ukraine*)
Matsumoto K.
(*Yamagata, Japan*)
Mormul P.
(*Warsaw, Poland*)
Mykhailyuik V.
(*Chernivtsi, Ukraine*)
Plachta L.
(*Krakov, Poland*)
Pokas S.
(*Odesa, Ukraine*)
Polulyakh E.
(*Kyiv, Ukraine*)
Sabitov I.
(*Moscow, Russia*)
Savchenko A.
(*Kherson, Ukraine*)
Sergeeva A.
(*Odesa, Ukraine*)
Shelekhov A.
(*Tver, Russia*)
Volkov V.
(*Odesa, Ukraine*)
Zarichnyi M.
(*Lviv, Ukraine*)

Administrative committee

- Egorov B., chairman, rector of the ONAFT;
- Povarova N., deputy chairman, Pro-rector for scientific work of the ONAFT;
- Mardar M., Pro-rector for scientific-pedagogical work and international communications of the ONAFT;
- Fedosov S., Director of the International Cooperation Center of the ONAFT;
- Kotlik S., Director of the P.M. Platonov Educational-scientific institute of computer systems and technologies "Industry 4.0";
- Svytyy I., Dean of the Faculty of Computer Systems and Automation.

Organizing commitee

Kirillov V.
Konovenko N.
Fedchenko Yu.
Maksymenko S.
Cherevko Ye.
Osadchuk E.
Prus A.

Trace Regularization Problem On a Banach Space

Erdal Gül
(Yildiz Technical University, Mathematics Department, Istanbul, Turkey)
E-mail: gul@yildiz.edu.tr

Let $\mathcal{H}$ be a separable Hilbert space and let $S_1[\mathcal{H}]$ be the trace class operators on $\mathcal{H}$ (First Schatten Class, [3]).

Consider $\mathcal{H}_1 = L^2(\mathcal{H};\ [0,\pi])$ and define an inner product on $\mathcal{H}_1$ by:

$$(f,g)_{\mathcal{H}_1} = \int_0^\pi (f(t),g(t))_{\mathcal{H}} dt$$

for all $f, g \in \mathcal{H}_1$.

- With this inner product, $\mathcal{H}_1$ is also a separable Hilbert space.

Here, we study the same problem in [2], with $\mathcal{H}$ replaced by a arbitrary separable Banach space $\mathcal{B}$, under the following conditions:

(1) $Q(t)$ has a weak second-order derivative in $[0,\pi]$ and for $t \in [0,\pi]$, $Q^{(i)}(t)$ $(i = 0,1,2)$ is a self adjoint trace class operator on $\mathcal{B}$.
(2) $\|Q\|_{\mathcal{H}_1} < 1$.
(3) $\mathcal{H}_1$ has an o.n.b. $\{\varphi_n\}_{n=1}^\infty$ such that $\sum_{n=1}^\infty \|Q\varphi_n\|_{\mathcal{H}_1} < \infty$.
(4) $\left\|Q^i(t)\right\|_{S_1[\mathcal{B}]}$ $(i = 0,1,2)$ is a bounded measurable function on $[0,\pi]$.

It is clear from (4), that this is a nontrivial problem since, among other things, in the standard approach, there are a number of possible definitions of $S_1[\mathcal{B}]$ (see [3] and Pietsch [9]).
We assume that $\mathcal{B}$ is a continuous dense embedding in a separable Hilbert space $\mathcal{H}$ and for each $f, g \in \mathcal{B}$, $(f,g)_h = (f,g)_{\mathcal{H}}$ is the Hilbert functional on $\mathcal{B}$.

Theorem 1 (Polar Representation Theorem)**.** *Let $\mathcal{B}$ be a separable Banach space. If $A \in \mathbb{C}[\mathcal{B}]$, then there exists a partial isometry U and a self-adjoint operator T, with $D(T) = D(A)$ and $A = UT$. Furthermore, $T = [A^*A]^{1/2}$ in a well-defined sense.*

Def. If $A \in \mathbb{S}_1[\mathcal{B}]$, we called it a trace class (or nuclear) operator on $\mathcal{B}$.
* Since $\mathbb{S}_p[\mathcal{H}]$ is a two sided *ideal, it follows that the same is true for $\mathbb{S}_p[\mathcal{B}]$.
* For $1 \le p < \infty$, $A \in \mathbb{S}_p[\mathcal{B}]$ and $B \in \mathcal{L}[\mathcal{B}]$ then $AB, BA \in \mathbb{S}_p[\mathcal{B}]$ and

$$\|AB\|_{\mathbb{S}_p[\mathcal{B}]} \le \|B\|_{\mathcal{L}[\mathcal{B}]} \|A\|_{\mathbb{S}_p[\mathcal{B}]}$$

$$\|BA\|_{\mathbb{S}_p[\mathcal{B}]} \le \|B\|_{\mathcal{L}[\mathcal{B}]} \|A\|_{\mathbb{S}_p[\mathcal{B}]}$$

Lemma 2. *If $\lambda \notin \sigma(L_0)$ then $QR_0(\lambda) \in \mathbb{S}_1[\mathcal{H}_1]$*

Lemma 3. *The operator valued function $R(\lambda) - R_0(\lambda)$ is analytic in $\rho(L)$, the resolvent set of L, with respect to the $\mathbb{S}_1[\mathcal{H}_1]$ norm.*

Theorem 4. *The regularized trace formula for operator L on B with the conditions on operator function $Q(t)$ is given by*

$$\sum_{m=0}^{\infty}\left[\sum_{n=1}^{\infty}(\lambda_{mn} - \mu_m) - \frac{1}{\pi}\int_0^\pi trQ(t)dt\right] = \frac{1}{4}[tr(Q(0) + trQ(\pi)$$

References

[1] A. Grothendieck, *Products tensoriels topologiques et espaces nucleaires*, Memoirs of the American Mathematical Society, **16** (1955).

[2] E. Gül, On the regularized trace of a second order differential operator, *Applied Mathematics and Computation* 198: 471-480, 2008.

[3] T. L. Gill and W. W. Zachary, *Functional Analysis and the Feynman operator Calculus*, Springer, New York, (2016).

[4] J. Kuelbs, *Gaussian measures on a Banach space,* Journal of Functional Analysis **5** (1970), 354–367.

[5] P. D. Lax, *Symmetrizable linear tranformations,* Comm. Pure Appl. Math. **7** (1954), 633-647.

[6] G. Lumer, Spectral operators, Hermitian operators and bounded groups, *Acta. Sci. Math.* (Szeged) **25** (1964), 75-85.

[7] L. A. Lusternik and V. J. Sobolev, Elements of functional analysis, (English Translation) Fredrich Ungar, New York, (1979).

[8] G. Maksudov, M. Bayramoglu and E. E. Adıgüzelov, *On regularized trace of Sturm-Liouville operator on a finite interval with the unbounded operator coefficient*, Dokl. Akad. Nauk SSSR **30(1)**, (1984), 169-173.

[9] A. Pietsch, *History of Banach Spaces and Operator Theory*, Birkhäuser, Boston, (2007).

[10] S. J. Szarek, *Banach space without a basis which has the bounded approximation property*, Acta Math. **159**, (1987), 81-98.