International scientific conference

«Algebraic and geometric methods of analysis»

Book of abstracts



May 30 - June 4, 2018,
Odesa,
Ukraine

https://www.imath.kiev.ua/~topology/conf/agma2018

List of topics

- Algebraic methods in geometry
- Differential geometry in the large
- Geometry and topology of differentiable manifolds
- General and algebraic topology
- Dynamical systems and their applications
- Geometric problems in mathematical analysis
- Geometric and topological methods in natural sciences
- History and methodology of teaching in mathematics

Organizers

- The Ministry of Education and Science of Ukraine
- Odesa National Academy of Food Technologies
- The Institute of Mathematics of the National Academy of Sciences of Ukraine
- Taras Shevchenko National University of Kyiv
- The International Geometry Center

Program Committee

Chairman: Prishlyak A.
(*Kyiv, Ukraine*)
Balan V.
(*Bucharest, Romania*)
Banakh T.
(*Lviv, Ukraine*)
Fedchenko Yu.
(*Odesa, Ukraine*)
Fomenko A.
(*Moscow, Russia*)
Fomenko V.
(*Taganrog, Russia*)
Glushkov A.
(*Odesa, Ukraine*)
Haddad M.
(*Wadi al-Nasara, Syria*)
Herega A.
(*Odesa, Ukraine*)
Khruslov E.
(*Kharkiv, Ukraine*)
Kirichenko V.
(*Moscow, Russia*)
Kirillov V.
(*Odesa, Ukraine*)
Konovenko N.
(*Odesa, Ukraine*)
Maksymenko S.
(*Kyiv, Ukraine*)
Matsumoto K.
(*Yamagata, Japan*)
Mykytyuk I.
(*Lviv, Ukraine*)
Mikesh J.
(*Olomouc, Czech Republic*)
Mormul P.
(*Warsaw, Poland*)
Moskaliuk S.
(*Wien, Austri*)
Panzhenskiy V.
(*Penza, Russia*)
Pastur L.
(*Kharkiv, Ukraine*)
Plachta L.
(*Krakov, Poland*)
Pokas S.
(*Odesa, Ukraine*)
Polulyakh E.
(*Kyiv, Ukraine*)
Sabitov I.
(*Moscow, Russia*)
Savchenko A.
(*Kherson, Ukraine*)
Sergeeva A.
(*Odesa, Ukraine*)
Strikha M.
(*Kyiv, Ukraine*)
Shvets V.
(*Odesa, Ukraine*)
Shelekhov A.
(*Tver, Russia*)
Shurygin V.
(*Kazan, Russia*)
Vlasenko I.
(*Kyiv, Ukraine*)
Volkov V.
(*Odesa, Ukraine*)
Zadorozhnyj V.
(*Odesa, Ukraine*)
Zarichnyi M.
(*Lviv, Ukraine*)

Administrative committee

- Egorov B., chairman, rector of the ONAFT;
- Povarova N., deputy chairman, Pro-rector for scientific work of the ONAFT;
- Mardar M., Pro-rector for scientific-pedagogical work and international communications of the ONAFT;
- Fedosov S., Director of the International Cooperation Center of the ONAFT;
- Kotlik S., Director of the P.M. Platonov Educational-scientific institute of computer systems and technologies "Industry 4.0";
- Svytyy I., Dean of the Faculty of Computer Systems and Automation.

Organizing commitee

Kirillov V.
Konovenko N.
Fedchenko Yu.
Hladysh B.
Osadchuk E.
Maksymenko S.
Khudenko N.
Cherevko E.

On The Second Regularized Trace Formula for a Differential Operator with Unbounded Coefficients

Erdal GÜL
(Yildiz Technical University, Mathematics Department, Istanbul, TURKEY)
E-mail: gul@yildiz.edu.tr

Let H be an infinite dimensional separable Hilbert space. Let denote the inner product and the norm in H by $(.,.)$ and $\|.\|$, respectively and denote the set of all kernel operators from H to H by $\sigma_1(H)$. Let $H_1 = L_2([0,\pi];H)$ be the set of all strongly measurable functions f defined on $[0,\pi]$ with their values in H such that for every $g \in H$ the scalar function $(f(x),g)$ is measurable in the interval $[0,\pi]$ and

$$\int_0^\pi \|f(x)\|^2 dx < \infty.$$

In $H_1 = L_2([0,\pi];H)$ we consider the operators

$$L = L_0 + Q, \;\; L_0 = y^{'v} + Ay$$

with the same boundary conditions $y^{'}(0) = y^{'}(\pi) = y^{'''}(0) = y^{'''}(\pi) = 0$. Here the operator $A : D(A) \to H$ is a densely defined on H such that $A = A^* \geq I, \; A^{-1} \in \sigma_\infty(H)$ where I is identity operator on H, A^* is the adjoint operator of A and $\sigma_\infty(H)$ is the set of all compact operators from H to H. And, $Q(x)$ is an operator function satisfying the following conditions:

(a) $Q(x) : H \to H$ is a self-adjoint operator for every $x \in [0,\pi]$.
(b) $Q(x)$ is weakly measurable in the interval $[0,\pi]$ and for every $f, g \in H$, the scalar function $(Q(x)f, \; g)$ is measurable on $[0,\pi]$.
(c) The function $\|Q(x)\|$ is bounded on $[0,\pi]$.

Let $\gamma_1 \leq \gamma_2 \leq \cdots \leq \gamma_n \leq \ldots$ be the eigenvalues of the operator A and $\varphi_1, \varphi_2, \ldots, \varphi_n, \ldots$ be the orthonormal eigenvectors corresponding to these eigenvalues. Here, each eigenvalue is represented as many times as its multiplicity. Moreover, let the eigenvalues of the operator L_0 and L be $\mu_1 \leq \; \mu_2 \leq \cdots \leq \mu_n \leq \ldots$ and $\lambda_1 \; \leq \lambda_2 \leq \cdots \leq \lambda_n \leq \ldots$, respectively.

Lemma 1. *If $\gamma_j \sim a.j^\alpha \; (a > 0, \alpha < \infty)$ as $j \to \infty$ then the asymptotic formula*

$$\lambda_n, \; \mu_n \sim \; dn^{\frac{4\alpha}{4+\alpha}} \; as \; n \to \infty \tag{1}$$

holds where d is a constant.

Let $R_\lambda^0 = (L_0 - \lambda I)^{-1}$, $R_\lambda = (L - \lambda I)^{-1}$ be the resolvents of the operators L_0 and L , respectively. By the well known equality

$$R_\lambda = R_\lambda^0 - R_\lambda Q R_\lambda^0 \;\; (\lambda \in \rho(L) \cap \rho(L_0))$$

we have:

Lemma 2.

$$\sum_{q=1}^{n_p} ({\lambda_q}^2 - {\mu_q}^2 \;) = \sum_{j=1}^{s} M_{pj} + M_p^{(s)}$$

where

$$M_{pj} = \frac{(-1)^j}{\pi i j} \int_{|\lambda|=b_p} \lambda \; tr[(QR_\lambda^0)^j] d\lambda \;\; (j = 1, \; 2, \ldots) \tag{2}$$

$$M_p^{(s)} = \frac{(-1)^s}{2\pi i}\int_{|\lambda|=b_p} \lambda^2 tr[R_\lambda (QR_\lambda^0)^{s+1}]d\lambda. \tag{3}$$

Theorem 3. *If the operator function $Q(x)$ satisfies the conditions (a), (b), (c) and $\gamma_j \sim aj^\alpha$ $(a > 0, \alpha > \frac{8}{7}(3+\sqrt{2}))$ as $j \to \infty$ then ,*

$$\lim_{p\to\infty} M_{pj} = 0 \quad (j = 2, 3, 4, \dots).$$

Theorem 4. *If the operator function $Q(x)$ satisfies the following conditions*

i) *$Q(x)$ has weak derative of the 8-th order in the interval $[0, \pi]$ and the function $(Q^{(8)}(x)u, v)$ is continuos for every $u, v \in H$.*

ii) *For every $x \in [0, \pi]$, $Q^{(i)}(x) : H \to H$ $(i = 0, 1, ..., 8)$ are self-adjoint operators.*

iii) *For every $x \in [0, \pi]$, $Q^{(8)}(x)$, $AQ^{(2i)}(x) \in \sigma_1(H)$ $(i = 0, 1, ..., 8)$ and the functions $\|Q^{(8)}(x)\|_{\sigma_1(H)}$, $\|AQ^{(2i)}(x)\|_{\sigma_1(H)}$ $(i = 0, 1, ..., 8)$ are bounded and measurable in the interval $[0, \pi]$.*

and if $\gamma_j \sim aj^\alpha$ $(a > 0, \alpha > \frac{8}{7}(3+\sqrt{2}))$ as $j \to \infty$ then the formula

$$\lim_{p\to\infty} \sum_{q=1}^{n_p} [{\lambda_q}^2 - {\mu_q}^2 - \frac{2}{\pi}\mu_q \int_0^\pi (Q(x)\varphi_{j_q}, \varphi_{j_q})dx]$$

$$= \frac{1}{2}[trAQ(0) + trAQ(\pi)] + \frac{1}{32}[trQ^{(4)}(0) + trQ^{(4)}(\pi)] - \frac{1}{\pi}\int_0^\pi trAQ(x)dx \tag{4}$$

is satisfied. Here j_1, j_2, ... are natural numbers.

References

[1] E. Adıgüzelov, P. Kanar, The second regularized trace of a second order differential operator with unbounded operator coefficient,*International Journal of Pure and Applied Mathematics* 22(3): 349-365, 2005.

[2] E. Adıgüzelov, Y. Sezer, The regularized trace of a self adjoint dferantial operator of higer order with unbounded operator coefficient, *Applied Mathematics and Computation* 218: 2113-2121, 2011.

[3] RZ. Chalilova, On regularization of the trace of the Sturm-Liouville operator equation, *Funks. Analiz, teoriya funksiy i ik pril Mahaçkala* 3: 154-161, 1976.

[4] Gohberg IC, Krein MG, *Introduction to the Theory of Linear Non-self Adjoint Operators*, volume 18 of *Translation of Mathematical Monographs*. (AMS, Providence, RI), 1969.

[5] E. Gül, A regularized trace formula for differential operator of second order with unbounded operator coefficients given in a finite interval, *International Journal of Pure and Applied Mathematics* 32(2): 225-244, 2006.

[6] E. Gül, On the regularized trace of a second order differential operator, *Applied Mathematics and Computation* 198: 471-480, 2008.